

| | |
|---|--|
| <ul style="list-style-type: none">○ O2 need ≤ 1lpm○ Day 3 or more of illness○ Normal CXR, if done (no atelectasis or pneumonia) | <p>airway issues</p> <ul style="list-style-type: none">○ Worsening respiratory status or oxygen need |
|---|--|

| <u>Diagnoses</u> | <u>Contraindications</u> |
|--|---|
| <p><u>Medical</u></p> <ul style="list-style-type: none"> • Croup <ul style="list-style-type: none"> ○ Patients who received 2 racemic epi doses may be admitted to obs but should be observed for a minimum of 12 hours after the 2nd dose • Dehydration <ul style="list-style-type: none"> ○ Gastroenteritis ○ Diarrhea • Cellulitis/Hand or Foot Infection • Peritonsillar Abscess/Pharyngitis <ul style="list-style-type: none"> - Evaluated by ENT if suspected abscess • Toddler Ingestions <ul style="list-style-type: none"> ○ 12 months to 5 years of age ○ Normal mental status after 2 hours ED Observation ○ Ingested half-life less than 8 hours ○ Normal vital signs for age | <p><u>Medical</u></p> <ul style="list-style-type: none"> • Croup <ul style="list-style-type: none"> ○ Stridor at rest ○ Oxygen need ○ Accessory muscle use ○ More than 2 racemic epi doses • Dehydration <ul style="list-style-type: none"> ○ Abdominal pain ○ Vomiting without diarrhea ○ Na \leq 130 or \geq 150 if done ○ Hypoglycemic ○ Bloody stools • Cellulitis/Hand or Foot Infection <ul style="list-style-type: none"> ○ Joint involvement ○ Diabetes or immune compromised • Peritonsillar Abscess/Pharyngitis <ul style="list-style-type: none"> ○ Airway concerns • Toddler Ingestions <ul style="list-style-type: none"> - Seizure or arrhythmia possible effect - Ingestion of oral hypoglycemics, anti-hypertensives, CCB, beta- |

Ortho/Trauma

- Fracture – Rule-out Compartment Syndrome
 - Orthopedics involved

Diagnoses

Ortho/Trauma

- Stable Thoracic/Lumbar Fracture
 - Awaiting brace, spine consulted
 - Normal neurological exam

blockers, etc.

- Oxygen need

Ortho/Trauma

- Fracture – Rule-out Compartment Syndrome
 - Other injuries except lacerations/abrasions
 - Altered mental status
 - Abnormal neurological exam

Contraindications

Ortho/Trauma

- Stable Thoracic/Lumbar Fracture
 - Other injuries except lacerations/abrasions
 - Altered mental status